Steve Roy

COMMISSION
AI DiPietro, Chairman
William Neilsen, Vice Chairman
Virginia Winn
Mark Koppelkam
Richard Frothingham
Thomas Racine
Ellen Leff

DIRECTOR OF PUBLIC WORKS
George Crombie

ENGINEERING DIVISION
Steven Goodkind, P.E., City Engineer
Ann Daughaday, CME

FINANCE MANAGER Patrick J. Buteau



Working together for Burlington

STREET DIVISION Bryan K. Osborne

TRAFFIC DIVISION
Don Moriey

INSPECTION SERVICES DIVISION Scott Corse

EQUIPMENT MAINTENANCE DIVISIONBruce Maysilles

WASTEWATER & SOLID WASTE DIV. Thomas E. Moreau

WATER DIVISION Laurie Adams

CONSTRUCTION MANAGEMENT Jerry Tomlinson





1990 WATER QUALITY REPORT

SUMMARY OF RESULTS

FOR THE

BEACH/SWIM SEASON







SUMMARY OF WATER QUALITY RESULTS FOR THE 1990 BEACH/SWIM SEASON

INTRODUCTION

This document serves as a summary of the data compiled during the 1990 beach season. In accordance with this year's water quality monitoring program (revision B, June 1990), it contains bacteriological results of Burlington's public beaches plus the streams which discharge into the vicinity of these beaches, including Englesby Ravine. Data has also been collected on the Winooski River both upstream and downstream of Burlington's East and North Wastewater Treatment Plants.

BACTERIOLOGICAL TESTS

The types of bacteriological tests performed vary for the first time this year. In addition to the standard Fecal Coliform test, certain samples were run for Fecal Streptococcus and Escherichia Coli (E. Coli) bacteria. These two strains of bacteria are actual forms of pathogenic organisms instead of Fecal Coliform, which merely serves as an indicator organism for pathogenic material. The reasons for trying these strains of bacteria are twofold: (1) A 1984 EPA document titled "Health Effects Criteria for Fresh Recreational Waters" found a correlation between E. Coli concentrations and gastrointestinal illnesses in exposed swimmers whereas there was no correlation using Fecal Coliform results. (2) A Fecal Coliform to Fecal Streptococcus (FC/FS) ratio was completed on stream samples to try and determine if the contamination was due to human or animal activity.

REPORT OBJECTIVE

The volume of data presented here in summary sheets makes complex analyses difficult, however we will try to answer the following questions:

- Are high bacteriological results on the Winooski River being caused by other sources or Burlington's wastewater plants.
- Is the contamination of streams discharging into the vicinity of Burlington's beaches coming from human or non-human origin.
- Can we determine the approximate location of stream contamination.
- Are high beach counts and subsequent closings the result of these streams, combined sewer overflows or a combination of both.

- Is E. Coli or Fecal Coliform the most accurate and repeatable bacteriological test to perform.
- Should we use the State of Vermont's E. Coli limit for wastewater effluent on beach samples or should we adopt a different limit based on the above mentioned EPA report.

WINOOSKI RIVER SAMPLES

Samples were taken both upstream and downstream of North and East treatment plants on the Winooski River which were then run for fecal coliform. At the same time, effluent samples from the plants were also run. In order to separate the data into usable information, dry weather and wet weather conditions are summarized below. Dry weather is set as rainfall less than or equal to 0.1 inches of rain within 2 days prior to the samples being taken. Wet weather is defined as more than 0.1 inches of rain within 2 days prior to the samples being taken. This definition of dry versus wet weather has been based upon elevated levels of bacteriological contamination in the Winooski River. A summary of results is shown below:

FECAL COLIFORM RESULTS (colonies per 100 mls of sample)

DRY WEATHER (rain North Plant	ifall≤0.1 " with ·	in 2 days prior)	•
Upstream	122 ave	300 high	30 low
Downstream	91 ave	190 high	30 low
East Plant			
Upstream	181 ave	360 high	30 low
Downstream	36 ave	140 high	<1 low
WET WEATHER (rail	nfall > 0.1 " with	nin 2 days prior)	
North Plant			
Upstream	1318 ave	2400 high	240 low
Downstream	1098 ave	1860 high	280 low
East Plant			
Upstream	788 ave	1880 high	130 low
Downstream	893 ave	2060 high	<1 low

As expected, the wet weather counts are invariably higher than dry weather. This shows that stormwater runoff into the Winooski is indeed contributing to bacteriological contamination. During this sampling period the fecal coliform counts from both plants averaged one colony with a high

of four colonies. With the exception of wet weather data for East plant, the downstream counts were consistently lower than upstream. This is caused by the dilution effects of effluent with low coliform counts and the die-off of bacteria, to a smaller extent.

According to East plant personnel, there is no stormwater outfall between the plant's discharge point and the downstream sampling point which could add contamination. For some reason, two out of four wet weather samples had larger downstream values than upstream, however this is no cause for concern.

BEACH RESULTS

Beach samples were run for Fecal Coliform from North, Leddy and Oakledge beaches between May 31st and September 7th. The number of sampling times is different for each beach because of repeat samples after a violation occurred and the fact that North beach was open longer than the others. The results for each beach are summarized below with summary sheets and graphs for the months of June, July and August attached in Appendix A at the end of this report. Also, during the month of August, all beaches were tested for the presence of E. Coli. Later on we will try to attribute the source of high coliform counts.

North Beach

Fecal Coliform

Forty-two (42) samples were taken from the north and south ends of North beach between May 31st and September 7th.

The north end had 7 violations of the Fecal Coliform limit out of 42 (17% violation) with only one that couldn't be attributed to a rain event as defined earlier. The violations ranged from 240 to 1700 colonies. The south end had 12 violations (29%) with three that again could not be attributed to rain events. High counts ranged from 220 to 1000 colonies.

E. Coli

Nine (9) samples were run for E. Coli between August 2nd and August 17th. The north and south ends had 4 violations of the 77 col/100 ml limit, all attributed to wet weather.

Leddy Beach

Fecal Coliform

Twenty-two (22) samples were taken from the north and south ends of Leddy beach between June 18th and August 17th.

The north end had no violations. The south end had one violation which may have been caused by a rain event.

E. Coli

The north end had 3 and the south had 4 violations out of seven (7), each having a high count on Aug. 17th, when rain was recorded at Main plant but not at North plant.

Oakledge Beach

Fecal Coliform

Twenty-three (23) samples were taken from the north, south and cove of this beach between June 18th and August 15th.

The north end had 7 violations (30%) with one not caused by a rain event. The south end had 5 violations (22%) with two not caused by precipitation. The cove had 4 violations (17%) with three that couldn't be attributed to a rain event.

E. Coli

The north, south and cove areas had 2, 1 and 1 violations respectively out of four (4) samples. All could be attributed to wet weather conditions.

COMBINED SEWER OVERFLOWS (CSO)

By definition, a CSO is said to have occurred when the stormwater flows are high enough to cause both storm and untreated wastewater to bypass the Main treatment plant and discharge directly into Lake Champlain in order to protect the plant from damage. Although it is difficult to know when stormwater contains raw sewage, it is likely that a CSO has occurred when the flow into Main plant has exceeded 7 MGD for more than 4 hours. Note, however, that these estimates did not always coincide with high total daily rainfall measurements at Main plant. CSOs are more a function of the intensity of a storm rather than the quantity that fell. For instance, a 0.25" recording in an hour can potentially be more devastating than 1" of rain over four hours. Using this assumption, 12 CSOs were recorded for the months of June, July and August of 1990. The dates are shown below:

Combined Sewer Overflows (Q > 7 MGD for > 4 hours) June 3, 21, 22, 23, 29 July 20, 23, 31 August 6, 10, 27, 28

This information will be used later to interpret beach results.

BEACH STREAMS

Streams feeding into each of these beaches were tested on a periodic

basis for Fecal Coliform (FC), Fecal Streptococcus (FS), and E. Coli (EC) and were found to be quite contaminated. There is one stream feeding into North beach, which has been designated as north stream. There are two streams feeding into Leddy beach, designated Leddy Stream North and South. Finally, the mouth of Englesby Ravine is located at the extreme north end of Oakledge beach. Samples of Englesby were taken at six locations from the head to the mouth. Also, a small stream located at the south end of Oakledge was sampled.

Fecal Coliform and E. Coli results are shown graphically in Appendix B at the end of this report. A log scale for the bacteriological counts was necessary because of the wide range of numbers. The last page of Appendix B shows EC versus FC plotted for all streams to see if there was any correlation between the two tests. All streams were grouped together in order to get over 30 data points, which statistics will tell you is the number of points necessary to make accurate asumptions. With a correlation coefficient of only 0.79, Burlington concurs with the EPA that there is no correlation between Fecal Coliform and E. Coli. Fecal Streptococcus will be used only as a ratio to determine the type of contamination.

A wet/dry weather scenario like the Winooski River samples was attempted, but since the watershed for each stream is different, this distinction was difficult and would have led to erroneous interpretations. For any distinct sample results mentioned, this author will point out any rainfall within two days prior to the sample being taken.

Summary of Stream Sample Results

North Beach Stream

Samples were taken from the head, middle and mouth of this stream. Results ranged from a low of 80 to a high of 13000 col/100 ml. Only 3 out of 21 samples were below the Fecal Coliform limit of 200 col/100 ml (86% above) and none were below the E. Coli limit of 77 (100 % above). Even in dry weather, this stream ran high counts. Comparing all three sample locations, the highest counts alternated fairly evenly between the head and mouth of this stream for FC, and the mouth was always higher than the others for EC. There does not appear to be a certain area causing contamination, since this data shows that the entire watershed is for the most part contributing.

Leddy Beach Streams

Eight FC samples and six EC each were run from the north and south streams at Leddy Beach. The north stream FC counts ranged from a high of 3800 colonies to a low of 20 with 5 out of 8 (63%) above the limit. North stream EC counts ranged from 50 to 1500 colonies, with 5 out of 6 (83%) above the limit.

South stream FC counts ranged from 20 to 5200 colonies, with 50% over the limit. EC ranged from <10 to >1000 with 83% over the limit of 77 col/100 ml.

North stream counts were higher than south stream for a given sample 5 out of 8 times which says that both streams are equally contaminated.

Oakledge South Stream

Eight FC and six EC samples were run for this stream. Fecal Coliform ranged from a low of 10 to a high of 4400 colonies with 88% over the limit. E. Coli samples ranged from 370 to 4400, and as you can see were all over the 77 col/100 ml limit. This stream appears to run contaminated all the time.

Englesby Ravine

Englesby Ravine was sampled in six locations, with some locations being done more often than others. For graphical purposes, three were designated Upper Englesby and three were called Lower Englesby. The lower section is comprised of the mouth, the ravine at Pine Street and Shelburne Road. Upper Englesby is the head of this ravine at Crescent Road and includes two ponds at Burlington Country Club which overflow into Crescent Road.

As expected, flows observed in the upper ravine area at all times were magnitudes of order less than the lower sections.

All results are summarized below, from head to mouth. The last two columns show the number of times counts were over the limit for a given parameter and how many times a particular location had the highest concentration compared to the others.

FECAL COLIFORM

Location	# sampling	# over	# highest for
	<u>times</u>	<u>limit</u>	sampling date
BCC Left Pond BCC Right Pond Crescent Road Shelburne Road Pine Street Mouth	7 7 7 8 5	5 (71%) 5 (71%) 7 (100%) 7 (88%) 5 (100%) 6 (67%)	1 0 1 3 1 3

E COLI

<u>Location</u>	# sampling <u>times</u>	# over <u>limit</u>	# highest for sampling date
BCC Left Pond	5	4 (80%)	1
BCC-Right Pond	5	3 (60%)	0
Crescent Road	4	4 (100%)	0
Shelburne Road	6	6 (88%)	2
Pine Street	5	4 (80%)	0
Mouth	8	6 (75%)	3

From the above data, one can see that this ravine like the others runs contaminated pretty much all of the time. High counts are less critical in terms of beach contamination at the head than at the mouth, since smaller flows means less of a total count of bacteria entering the lake. The source of contamination does not originate from one particular area, rather it starts at the head and is continually added throughout the length of this ravine. Reasoning for this hypothesis is such; if bacteria was being added only at the head of this ravine, increasing flows added by travelling downward toward the lake would reduce the concentration of bacteria per 100 mls of sample. One can see from this data that the counts typically increase as they travel towards the mouth which tells us that bacterial contamination is being added at a rate greater than the addition of increased flow. Appendix D shows a graphical representation of this hypothetical situation.

FC/FS RATIO

The Fecal Coliform to Fecal Streptococcus ratio has been employed to determine whether the source of bacterial contamination stems from human or non-human origin. According to the 17th Edition of "Standard Methods for the Examination of Water and Wastewater", a FC/FS ratio <0.7 indicates animal origin, 0.7 to 4.4 is mixed human and animal, and >4.4 is human only. This version of "Standard Methods" makes it clear that this ratio is only a guide and many factors can skew the results. One of the most critical factors is the relatively short lifespan of the streptococcus bacteria outside its host compared to fecal coliform. However, it is this author's belief that if the streptococcus had indeed died off quicker than the coliform bacteria, it would only skew the ratio toward the mixed human/animal area since FC is in the denominator of the equation. The results, given at the end of this report, are summarized here.

Leddy Stream North - 7 samples taken	6 animal	1 mixed
Leddy Stream South - 6 samples	5 animal	1 mixed

North Stream Mouth - 9 samples taken North Stream Middle - 2 samples North Stream Head - 5 samples	6 animal 2 animal 5 animal	3 mixed
Englesby Ravine Mouth - 7 samples taken ER @ Shelburne - 5 samples	4 animal 5 animal	3 mixed
ER @ Crescent - 5 samples	4 anima1	1 mixed
BCC, left pond - 5 samples	4 anima1	1 mixed
BCC, right pond - 6 samples	6 animal	
Oakledge Stream - 6 samples taken	5 animal	1 mixed

One can see that at no time did the ratio indicate only human contamination. As a check, the ratio was done on primary effluents of all three treatment plants twice to see what would happen. Although the extremely high dilutions necessary to read the plates can produce large errors, the ratios ranged from a mixed of 0.8 to a definitely human ratio of 27.7.

In August of this year, a large portion of Englesby Ravine was walked to find any evidence of human contamination. No pipes were found illegally discharging into the ravine, and no evidence was found along the banks. Since the ravine and other monitored streams drain large amounts of land, both urban and wooded, this author believes that animal contamination of the nature found is a very real problem.

CSO VERSUS STREAM

This next section is an attempt to determine if high Fecal Coliform counts were perhaps caused either by a CSO or a stream. The dates of beach violations are shown below with potential causes.

Beach <u>Location</u>	Violation <u>Date</u>	Main Plant <u>CSO</u>	Beach <u>Stream</u>	<u>Comments</u>
North (South end)	18 June	No	Unknown	Stream samples started on 27 June
Oakledge (All)	20 June	No	Potential	Probably Englesby; 0.04 " rain fell on 19 June
Oakledge (Cove)	25 June	No	Potential	Maybe caused by Oakledge South Stream
Oakledge (All)	28 June	No	Potential Potential	Colonies seen in Englesby

			. •		
rr f.	,				
	Oakledge (All)	5 July	No	Probable	0.05" rain fell on 4 July 🗸
	Oakledge (South)	6 July	No	Potential	Next sampling not until 10 July
	North (A11)	9 July	No	Probable	Background levels seen on 10 July
	North (South)	12 July	No	Probable	Background levels seen on 11 July
•	Oakledge (No. & So	16 July a.)	No	Probable	High levels seen on same day
	North (South)	23 July	No	Probable	High levels seen on same day
	North (All)	24 July	Potential	Potential	CSO on 23 July plus high stream levels
	Oakledge (North)	24 July	Potential	Potential	High levels in Englesby
	North (A11)	6 August	Potential ·	Probable	CSO same day; very high levels in stream
	North (South)	7 August	Potential	Probable	Yery high levels in stream
	North (All)	9 August	No	Probable	Very high levels seen on 7 August
	Oakledge (North)	13 August	No	Probable	High counts throughout Englesby
	North (All)	15 August ,	No	Potential	Reinfall on 13 (0.5") and 15 (0.05") August
	Leddy (North)	15 August	No	Potential	Rainfall on 13 (0.65") at North Plant
	Oakledge (North)	15 August	No	Potential	Rain plus high levels seen in Englesby on 13th
	North (A11)	28 August	Potential	Potential	CSO on 27 August; no samples taken after 20 August
The state of the s	North (All)	29 August	Potential	Potential	CSO on 28 August
	North	7 Sept.	Unknown	Unknown 9	Data not available

This information shows that the streams feeding into our beaches are much more of a threat than a CSO from the Main Treatment Plant. Therefore, combined sewer separation and stormwater treatment may reduce but <u>will not eliminate</u> the number of beach closings due to bacteriological contamination. The only options available to reduce loadings on the beaches caused by contaminated streams are briefy outlined below:

Option	<u>Advantage</u>	<u>Disadvantage</u>
Source elimination or reduction	Ideally the best way to clean up streams	Nearly impossible since source does not originate from a particular area. Non-enforceable
Detention Ponds	Contaminated stream flow not allowed into Lake	Not possible on all streams because many are too large
Stream Disinfection	Bacteriological kill achieved over wide flow range	Disinfection by-products released if chlorine used. Continual 0 & M costs

FECAL COLIFORM VERSUS E. COLI

This last section deals with the use of one bacteriological test over another for next year's beach/swim season and beyond. Fecal Coliform is still widely used for recreational waters, and in fact will continue to be used by the State of Vermont Health Department. However, a 1984 EPA report titled "Health Effects Criteria for Fresh Recreational Waters" could not find a correlation between FC counts and gastrointestinal illnesses in exposed swimmers. Although this report is the only one known to us, it is quite thorough in its procedure. EPA did find a correlation between illnesses and concentrations of both E. Coli and Fecal Streptococcus. We have successfully tested samples for streptococcus and the procedure is easier than E. Coli, however its 48 hour incubation period makes it unuseable for beach testing. Our next choice is between using FC or EC as a standard, and at what concentration is considered an acceptable risk. Let's review the current standards and use them in conjunction with our own data to make this determination.

The current wastewater standard for Fecal Coliform is 200 colonies per 100 mls of sample. This standard is derived from the Total Coliform limit, which was arbitrarily set by the California Bureau of Sanitary Engineering in 1943 at 1000 col/100ml. The FC limit was then adopted after research in the mid 1960's showed that Fecal Coliform averaged roughly 20% of Total Coliform in the Ohio River. Therefore, neither of these standards have any epidemiological basis.

Based upon the above mentioned report, EPA has asked States to utilize E. Coli as the standard for wastewater effluent. They gave States the flexibility to

set the limit, however they could not exceed 125 col/100 ml, which translates to a level of risk for 9 out of 1000 swimmers (0.9%) to potentially contract gastrointestinal illness by swimming in polluted waters. The State of Vermont has adopted a wastewater effluent limit of 77 col/100 ml which translates to 6 infected swimmers out of 1000 (0.6%). Appendix C contains the graph taken from this EPA report. Burlington does not need to adopt the State's effluent limit for recreational waters, but can make their own choice about an acceptable level of risk. A comparison of the beach violations using Fecal Coliform, E. Coli at 77 col/100 ml and E. Coli at 125 col/100 ml are summarized below:

<u>Violation Comparisons</u>

Beach <u>Location</u>	% violation Fecal Coliform	% violation E. Coli @ 77	% violation E. Coli @ 125
North Beach North	17%	44%	33%
North Beach South	29%	44%	22%
Leddy Beach North	0%	43%	14%
Leddy Beach South	5%	57%	29%
Oakledge North	30%	50%	50%
Oak ledge South	22%	25%	25%
Oak ledge Cove	17%	25%	0%

It is this author's belief that E. Coli be considered as the method of choice because of the epidemiological evidence presented by the EPA. In addition, I also think that the recreational water limit in Burlington be 125 colonies per 100 mls of sample because the level of risk at this concentration (0.9% versus 0.6%) is, in my opinion, acceptable and would not cause unnecessary beach closings. The wastewater limit of 77 colonies is acceptable and achievable for a water that is disinfected, but this limit at the beaches would result in overly conservative closings. Unfortunately, too few samples were taken this summer to immediately adopt this policy. Therefore, it is recommended that a decision be made after next summer's data is reviewed. Next summer's goal would be to run Fecal Coliform and E. Coli on the same samples, using the FC results as a governing factor.

SUMMARY OF OBJECTIVES


The objectives outlined on pages 1 and 2 of this report are summarized below:


- Are high bacteriological results on the Winooski River being caused by other sources or Burlington's wastewater plants.
 High Fecal Coliform counts on the Winooski River result from stormwater runoff during rain events and are not caused by Burlington's treatment plants.
- Is the contamination of streams discharging into the vicinity of Burlington's beaches coming from human or non-human origin.
 Evidence shows that the contamination of streams feeding into Burlington's beaches are caused with high certainty by animal sources.
- Can we determine the approximate location of stream contamination.
 Based upon the information presented, bacteriological contamination is added throughout the drainage area of these streams. No specific locations can be identified.
- Are high beach counts and subsequent closings the result of these streams, combined sewer overflows or a combination of both.
 Most of the beach closings appeared to be caused by streams flowing into them rather than CSO's. Out of all the beaches, North Beach appears to be the most sensitive to a potential of CSO's based on its geographical location. CSO locations at the time Main Plant, Maple Street, College Street
- Is E. Coli or Fecal Coliform the most accurate and repeatable bacteriological test to perform.
 E. Coli is the best method based upon scientific evidence.
- Should we use the State of Vermont's E. Coli limit for wastewater effluent on beach samples or should we adopt a different limit based on the above mentioned EPA report.
 It is recommended that we adopt a limit of 125 colonies/100 mls of sample for recreational waters after sufficient data from next year's beach/swim season is reviewed.


Many thanks to Burlington Parks Department for collecting beach samples, and especially to DPW's Wastewater Division, both full-time and part-time employees, who not only collected but also ran and read most of the samples done this summer.


Stephen T. Roy Process Engineer


APPENDIX A































APPENDIX B

























UPPER ENGLESBY RAVINE (E. COLI)



OAKLEDGE SOUTH STREAM (E. COLI)







Fecal Coliform

APPENDIX C

Taken from "Health Effects Criteria for Fresh Recreational Waters", Environmental Protection Agency, 1984.






Figure 6. Criterion for estimating the geometric mean *E. coli* density per 100 mL from an acceptable risk level of swimming-associated gastrointestinal illness.

APPENDIX D





RAW DATA

Sample:	Sample:	Sample:	fα(ν∼ Sample:	Zainnie:	Sample:	Sample:	Sample:	Sample:	. Sample:	Sample:	Sample:	Sample	eria () . Serie	edi. Si Sirik	en de en decidente .
· ·	e: 8.22		e: 8/13	30	[3]	e: 1.30	e: 7.35	e: 7.33	e: 1/6	%	10	667	SAMPLE DATE BY	"TESTING"	
	2) i pe.	13	7	-		- 3/2	 N	 	2.25	5.5	1.6.		.ING.	
		(2)		À,100.	300	120		3,800	710		240	110	LEDDY BEACH	_ *	ς.
		(30)		500	× 100	93d		720	40		410	.5,200.	Norll	Fecal * BURLINGTON	
		(2)		12,000	10,000.	2110	2,100	TNIC	900	130	180	3,400	NORTH BEACH MOUTH		
							1.900	3, 400	7,600	270			Mibore		
		1,600		13,000		1220	1500	2,400	11,500	230			HEAD	- WASTEWATER DIVISION	
	1/0		2200	7000	Of too	.660		(b,100)	5,400		60	40	nouth	/ISION	
	350		200	1,000		380		2,500	2,100		230	580	Shelburne		·
	320		400	7600		530		2,200	1,890.		750		ENGLESBY RAVINE (Atscent	Sheet #	
	290		Moge	8,000		530		4,600	HOF STORY			-350-	Seuth Prospert	1-08	
	310 90		300	4000		(3cy		1900 530	3,500 / 890		30 340		L. Gally R.		
	470	[saco	4000	600	040 240	:- (- 4,400	900	[ō	•	CAR leage South		

Sam	Sa	Sa ₁	Sa	Sa	က ည	Sa	Sa	Sa	Sa	Sa	ഗ്ദ			
	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:			X
01/22/90	09/22/90	6/21/90	6/21/90	6/20/90	6/19/90	6/18/90	6/14/90	411/90	6/7/90	6/4/90	5/31/90	SAMPLE DATE	Fecul "TESTING"	\
Parks	Porks	Parks	Parks	Parks	Parks	Parks	Parks	Parks	Parks	Parks	Parks	SAMPLED BY		
M	AM	PM	AM									LAB	BURLINGTON PUBLIC	
cab accident		-	2	*	20	14	22	48	48	126	170	NORTH BEACH	-	
Lab			Sign Land Chineses		50	360	92	2	20	22	14	BEACH SOUTH	WORKS - WAST SWIM BEACH	<u></u>
cab				20	4	20						 LEDDY E NORTH	WASTEWATER DIV	
Lab accident 1 c				8	Consideration .	24						BEACH SOUTH	DIVISION	
Lab accident	£10	34	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	800		4						NORTH		
accident	1-2/0-1	80	-1.40 idia 11476	1800	V (Treath V	^						OAKLEDGE SOUTH	Sheet #	
Lab t	1 / 1	6	76	390	<u>V</u>	2						COVE		9

"TESTING"

BURLINGTON PUBLIC WORKS - WASTEWATER DIVISION SWIM BEACH SAMPLES Sheet #

Sar e:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	
7/18/50	7/16/90	7/13/90	7/12/90	7/11/90	7/10/90	7/9/90	7/6/90	7/5/90	06/2/4	6/28/90	6/25/90	SAMPLE
Parks	Parks	Parks	Parks	Parks	Parks	Parks	Parks	Parks	Parks	Parks	Parks	SAMPLED BY
e de l'annue de préside me l'angle de l'annue de l'annu												LAB
	52-	/0	ω	/0	10	(200)		1/2		34	00	NORTH NORTH
	26	V0	1000	^	20	(254)		28		^	6.4.6	BEACH SOUTH
	30		2			98		43		2	6	LEDDY BEACH NORTH SOUTH
	118		2			30,444		20		2	/0	BEACH SOUTH
4	1400		/2			28	20	(700)	10	(600)	28	NORTH
0	(600)		4			4	(220)	7400	المقدمول على المقالمة المائية	500	08 in Administra	OAKLEDGE SOUTH
2	56		<u>v</u>			Hall Manufacture B	20	(600)	0-1	206	700	COVE

Sam 'e:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:	Sample:		
8-13-90	8-10-90	8.9.90	8.7.90	8.6.90	8.2.90	7.30.90	7.26.90	7/24/90	7/24/90	7/23/90	7/19/90	SAMPLE DATE	"TESTING"
PARKS	PARKS	PARKS	PARKS	PARKS	PARKS	PARKS	PAKKS	Parks	Parks	Parks	Parks	SAMPLED BY	
AM	P _M	I AM	PM	AM	AM	AM	M	PM	AM			LAB	BURLINGTO
-20	150	1,000	J/O	340	1/20	56	86	90	(240)	40	72	NORTI NORTH	BURLINGTON PUBLIC W
20	To Some	450	OBB	370 20 80	180	124	30	70.4	360	£20	90	NORTH BEACH RTH SOUTH	WORKS - WAS SWIM BEACH
10		100		1 80	20	30	16	20	20	70	18	LEDDY NORTH	- WASTEWATER DIVISION BEACH SAMPLES
160		140		20	40	122	18	100	20	V/O	98	BEACH SOUTH	IVISION
(610)		30		40	120	16	16	003/1200				NORTH	
/30		40 70			ನಿಂಂ	74	8	1 2/00m 130				OAKLEDGE SOUTH	Sheet #
8		70		50.3836 2000	30	4	8/	130				COVE	4

PARKS PM 30 30 30 30 PARKS PM 300 1000 1700 800 1000 1000 1000 1000 100	"TESTING" SAMPLE DATE Sample: 8.15.90
5 PM 300 100 770 80 30 30 30 30 30 30 30 30 30 30 30 30 30	
5 AM 30 -20 - 80 S	8-17-90 7
5 AM 30 400 5 AM 30 400	<u> </u>
5 AM 30 - 20 800 S	8.23.90
5 PM -20 -380 5 PM -30 -380 5 PM -30 -380	8-27-90 7
5 AM 30 -20 80 5 AM 30 -210 -210 -	1 06.38.8
5 AM 30 - 380 - 380 - 380 - 380 - 30	1 13.34.3
9 AM 1 30 m 1 30 m 1 30 m 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 08.86.8
J PM 1 510-1-	8-30-90 F
66	8-30-90 9
	9-7-90 2

¥
3
Col
·.

	1			77 (027								
	"TESTING"		BURLINGTO	BURLINGTON PUBLIC WORKS		- WASTEWATER DIVISION	VISION		Sheet #	90-1		
	SAMPLE !	LEDDY BEACH	BEACH	NORTH	STREAM	E.R. SAMPLES			WASSELVEN			
	ВУ	7	SOUTH	MOUTH	South Presp	Pasa Michigan	Mouth	conclude	FLANKANEN A	SMC A	1 Mark Smet	ナスト
Sample:	6/27	210	TNTC	~ /600	330				Coft	2194+		
Sample:	7/11 5.6			240								
					ਮੇਮੁੱਟ					***************************************		-
Sample:	7.23 S.F	1500	800	TUTC	2,400	2,570	(6100)	4400	1600	680	2,400	12,500
Sam la.	7.1C. SE			0707								
Complete.	1 03			B 140							1,000	1,000
Sample:	7. 30 St	160	110	1960	3/0	430	040	410	Joseph	60		- 840
	S .	ن ا		2.300								 1
0.000								6,6,5				
Sample:	8,7	1200	300	15,000	6000	3000	(good)	4000	6000	2000		3000
	8./2				5	200	-					1
Sample:					800	100	100	1600	100	100		
Sample:	8,20	50	入10				2/0					1440
Sample:	27.8				10	370	۸ ₁₀	370	1 /6	7,		1
			-									-
Sample:												
Specific.												1
) Salduste:												
Samole.	: >				· -							
, t											_[-

.

.

Sheet # 70-1

Sample:	Sample	Sample:	Sample	Sample:	Sample	Sample:	Sample	Sample:		Sample	Sarc	Sample	ording.	Sample:	
					8.15	8.8	8,0	7.36		7.33	7/10 76	7//8	, 1	7/, 7.6.	DATE
										880,000					Mary CSO
					2200	ಪಿಎಂ	50	01				30		200	NORTH
					400	0	10	370				50			MOUTH WR1
					/300	0	180	350				110			DOWNSTREAM NORTH WR2
					TNIZ							<u> </u>		· //	NORTH WATP EFFLUENT
					//00	٥	70	940				/30			UPSTREAM NORTH WR3
					700	0	160					10			DOWNSTREAM EAST WR4
					TRATE							\\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \		. </td <td>EAST WATP EFFLUENT</td>	EAST WATP EFFLUENT
					700	0ئو	100					310			UPSTREAM EAST WRS
[ı 		JAIN .				,		· l	1	'	!	Herry

	"TESTING"		BURLINGTON PUBLIC	-	WORKS - WAST	WASTEWATER DIVI	VISION		Sheet #	90-1
	SAMPLE DATE	SAMPLED BY	LAB	NORTH BEACH	H BEACH SOUTH	LEDDY NORTH	SOUTH	NORTH	OAKLEDGE SOUTH	COVE
			The statement of the st							
Sample:	8.2.90	PAKKS		40 40	day	0 -3) d	2.2	100	X/10
Sample:	8.6.90	PARKS		1/20 -	(NO)	5.7	70	40	10 wet	6. 8
Sample:	8.7.90	PARKS	B	30	v.et 20					. ,
Sample:	8.9.90	PARKS		190	1000	/0/ 1-ard	80 ,			
Sample:	8.10.90	PARKS		40	70					
Sample:	8-13-90	PARKS		/0	20 20	/0 /0	we.\ 2/50	330	130	108 TH
Sample:	8.15.90	2naks		370	330 330	500	080 + \$0	120	vet 60	00°
Sample:	8.17.90	PARKS	AM	30	. 8 tm	مريز)	in, 70			
Sample:	8-17-90	PAKKS	PM	: 1200	1300 1300	08.	300			
Sample:										
Sample:										
Samp : 1										

Fecal Coliform / Fecal Overprocession
BURLINGTON PUBLIC WORKS - WASTEWATER DIVISION
SAMPLES /Fecal Streptococcus Ratio

Streams into Beaches

	•		יייייייייייייייייייייייייייייייייייייי		747			
		6/27	7/10	7/16	7/11 .	7.23	7,25	7.30
Sample:	STREAM NOTA	21.0	1.26	0.607				0.20
Sample:	STEAM South	0.50	<0.1	0.005				1.43
Sample:	STLEAM MOTH	1.95	0.285	0.163	0.162		0,38	15%
Sample:	SICEMA MUNIC			0.355			0.29	
Sample:	STEETH SEXUE			0.326			0,33	0.45
Sample:	Englisby R.	0.15	0.25	0.386				18.0
Sample:	shelburg	0.29	<i>ن. ي</i> ي	0.07				0,53
Sample:	cresiont		0.53	0.03		0.19		0.76
Sample:	Englesby R.	0.37		-		0.27		0.15
Sample:	Left Calfland		25.0	0.08		0.10		181
Sample:	Eight Cell lase		0.07	0.07		0.05		0.04
Sample:	south Hall		0.04	0,103				0.41
Samr'a:								
				֡				

0.7-4.4: Mixed >4.4: HUMAN KO.7 : Animal

20.7: AND 0,7-4.4:11, Kid >4.4 : Home Samr 9: Sample: Sample: Sample: Sample: | Thest Sample: Crsunt Sample: | Shelburne Sample: Sample: Sample: LEDDY Sec Ay Sample: Sample: Scimple: LEDAY Neith Call by Gelf Right Neigh Mogh North Street HEAD CATREORY l EIR, "TESTING" SAMPLE DATE *ج*ہ کے, 27. 64 31 6,30 ヘ . . 0 09.0 0.82 1,43 80000 20.0 0,06 BURLINGTON PUBLIC WORKS - WASTEWATER DIVISION 0,22 0.16 0.36 Food Colfin / Front Stopt Ruhe 0.59 0.20 0.57 0,33 o.as وي دو 81.3 /:/ 0,008 6.0.2 2002 0,40 Sheet #

Fecal Colifera / Fecal Streptococcus Ratio
BURLINGTON PUBLIC WORKS - WASTEWATER DIVISION
SAMPLES

Swim Eweches, plus Primary

Sample: M	Sample:	Sample: 19.	Sample: IA	sample: 识别	Sample:	Sample: CA	Sample: _S	Sample: A	Sample:	Sample:	Sample: 1	Sample: 1 Mb	
MAIN (SO		Primery	Aff. North	EFF HAIN!		Coxe	South	North	Leddy South	Leddy Bosch North	NEATH GRAH	NOCTH N	
		Q. & 2	13.60	27.70							< ado	.]	7/10
		0,80 9.34	1 47 5 PW 1	16.37 185 69 14	ļ. 								7/16
2.95													7.23
						0.26	0.62	0.71	1.07	₹1.0	1.0	0.13	8.13
													·

20.7: AnimAL

>4.4 : HUMAN